

STAT⁺CUP[®] II



STATCUP[®] II

The STATCUP[®] II is a fully integrated on-site urine drug test cup. Simply collect the sample and read the results. It's that easy! See back for full procedure details and how to read results.

- Fast
- Accurate
- Reliable
- Easy-to-Use
- Affordable
- Dark Lines
- FDA Approved
- CLIA Waived
- Health Canada
- CE Marked
- ISO Certified Manufacturing

CALL FOR A
FREE SAMPLE!

TEST FEATURES

- No Urine Exposure
- No Steps to Activate
- Screw Top Lid
- Temperature Strip
- Test Cup Holds up to 18 Drugs
- Available with Built-in Adulterants
- Transport Sample to Lab

Drugs of Abuse:

Marijuana	Propoxyphene*
Cocaine	Fentanyl*
Opiates/Morphine	Tramadol*
Methamphetamine	K2/Spice*
Amphetamine	6-Monoacetylmorphine*
Phencyclidine	Cotinine*
Benzodiazepines	Alcohol*
Barbiturates	Ketamine*
Methadone	Bath Salts*
Ecstasy	Ethyl Glucuronide (ETG)*
Tricyclic Antidepressants	
Oxycodone	
Buprenorphine	
Methadone Metabolite	

Adulterants:

pH	Creatinine
Specific Gravity	Nitrites
Oxidants	Glutaraldehyde

*For forensic use only. Not intended for treatment or diagnostic purposes.



Toll Free: 1-800-668-0031

www.Tryrapid.com




STAT⁺ CUP II


**EASY-TO-USE
FAST
ACCURATE**

PROCEDURE



1 Remove Cap.




2 Donor provides specimen.



3 Replace the Cap securely. Peel off the blue label that covers the result window.



INTERPRETING RESULTS




Interpret adulterant test strip between 3-5 minutes. Refer to enclosed color chart for an accurate interpretation.



Phone: 1-800-668-0031

Rapid Response Drug Testing



NEGATIVE POSITIVE INVALID

All positive results are presumptive and should be confirmed by an alternative method (e.g. GC/MS). A photocopy of the results can be made by placing the entire sealed cup with the card face down on the copier.